

PRAHA2016

The 24th International Conference on High Resolution Molecular Spectroscopy

Prague, Czech Republic, August 30th-September 3rd, 2016

Výbor mezinárodní konference zve všechny zájemce na přednášku autorů
William E. Perreault, Nandini Mukherjee, and Richard N. Zare, kterou prosloví

profesor Richard N. Zare

Department of Chemistry, Stanford University

na téma

Angular and Internal-State Distributions of Photofragments Determined from Time-of-Flight Mass Spectrometry

Přednáška proběhne v rámci cyklu Ioannes Marcus Marciho přednášek a bude se konat
ve Velké zasedací síni pražského zastupitelstva na Pražské radnici na Mariánském
náměstí pod záštitou primátorky Prahy

ve čtvrtek 1. 9. 2016 od 16:30 hod.


Jako předřečník vystoupí Dr. Jon T. Hougen (NIST, Gaithersburg, USA)

Angular and Internal-State Distributions of Photofragments Determined from Time-of-Flight Mass Spectrometry

William E. Perreault, Nandini Mukherjee, and Richard N. Zare

Department of Chemistry, Stanford University, Stanford, California 94305 USA

The anisotropy parameter β for the angular distribution of the photoelectron and photoion in (2+1) resonance enhanced multiphoton ionization process of $\text{H}_2 \text{X } ^1\Sigma_g^+$ ($v=0, J=0$) molecules through the intermediate $\text{H}_2 \text{E,F } ^1\Sigma_g^+$ ($v \neq 0, J \neq 0$) level ($\lambda = 201.684 \text{ nm}$) is determined without using imaging hardware. Instead, we record the arrival time of the H_2^+ ions using a time-of-flight mass spectrometer when a molecular beam of H_2 is photoionized. Time-of-flight spectra were obtained as the direction of polarization of the ionizing laser was varied with respect to the flight axis and were fitted to an angular distribution in an appropriately rotated coordinate system with the z -axis oriented along the time-of-flight axis. The anisotropy parameter β was found to be 1.72 ± 0.13 by fitting the time-of-flight spectra and agreed with previous measurements. Using secondary ionization with a delayed laser pulse of different wavelength we also determined the vibrational energy distribution of the ions, showing greater than 99% of the ions are generated in their ground vibrational state ($v^+=0$), in agreement with the calculated Franck-Condon factors between the $\text{H}_2 \text{E,F } ^1\Sigma_g^+$ ($v \neq 0$) and $\text{H}_2^+ \text{X } ^2\Sigma_g^+$ (v^+). The time-of-flight spectrometry technique presented here has general applicability for photofragmentation processes involving light molecules.



Time-of-flight spectra of the H_2^+ ion for (a) $\theta = 0^\circ$; (b) $\theta = 30^\circ$; (c) $\theta = 60^\circ$; and (d) $\theta = 90^\circ$. The green curves are fits to the blue data points, based on the fitting for $\theta = 0^\circ$. The angle θ describes the angles between the optical field of the laser and the flight axis of the molecular beam.

Support of the U.S. Army Research Office under ARO Grant No. W911NF-13-1-0126 and MURI Grant No. C13J11493 is gratefully acknowledged.